

REVIEW ARTICLE OPEN



Culturally adapting internet- and mobile-based health promotion interventions might not be worth the effort: a systematic review and meta-analysis

Sumeyye Balci¹✉, Kerstin Spanhel², Lasse Bosse Sander² and Harald Baumeister¹

Health promotion interventions offer great potential in advocating a healthy lifestyle and the prevention of diseases. Some barriers to communicating health promotion to people of certain cultural groups might be overcome via the internet- and mobile-based interventions (IMI). This systematic review and meta-analysis aims to explore the effectiveness of culturally adapted IMI for health promotion interventions among culturally diverse populations. We systematically searched on Cochrane Central Register of Controlled Trials (CENTRAL), EbscoHost/MEDLINE, Ovid/Embase, EbscoHost/PsychINFO, and Web of Science databases in October 2020. Out of 9438 records, 13 randomized controlled trials (RCT) investigating culturally adapted health promotion IMI addressing healthy eating, physical activity, alcohol consumption, sexual health behavior, and smoking cessation included. From the included studies 10,747 participants were eligible. Culturally adapted IMI proved to be non-superior over active control conditions in short- ($g = 0.10$, [95% CI -0.19 to 0.40]) and long-term ($g = 0.20$, [95% CI -0.11 to 0.51]) in promoting health behavior. However, culturally adapted IMI for physical activity ($k = 3$, $N = 296$) compared to active controls yielded a beneficial effect in long-term ($g = 0.48$, [95% CI 0.25 to 0.71]). Adapting health promotion IMI to the cultural context of different cultural populations seems not yet to be recommendable given the substantial adaptation efforts necessary and the mostly non-significant findings. However, these findings need to be seen as preliminary given the limited number of included trials with varying methodological rigor and the partly substantial between-trial heterogeneity pointing in the direction of potentially useful culturally adapted IMI which now need to be disentangled from the less promising approaches.

PROSPERO registration number: 42020152939

npj Digital Medicine (2022)5:34; <https://doi.org/10.1038/s41746-022-00569-x>

INTRODUCTION

Health promotion interventions are an effective way of disease prevention and improving overall health^{1,2}. Well-established approaches focus amongst others on healthy eating, exercising, avoiding excessive use of alcohol, quitting smoking, and sexual health behaviors, such as condom use^{3,4}. Promoting these health behaviors on a large scale by using health promotion interventions might be a promising mean for reducing the burden of disease⁵⁻⁷. However, these intervention offers do not always reach or fit populations equally between and within countries⁸.

The reach of these interventions could be expanded globally via the internet⁹ using the Internet- and mobile-based interventions (IMI)¹⁰⁻¹⁴. IMI offers time and place flexibility, potential cost-effectiveness, and scalability without necessarily losing effectiveness¹⁵. Thereby, IMI could reach a diverse group of people including minorities and people of a cultural background that is not yet well covered by the established health care systems¹⁶⁻¹⁹. However, IMI also come with some substantial limitations, most prominently risk of low adherence and uptake^{20,21}. In order to tackle these issues, population-related factors such as the needs and expectations of the users should be taken into account¹⁵. Tailoring the intervention content and delivery method to the target group's culture can thus be a means for increasing engagement and effectiveness²².

Researchers in the field of health promotion are encouraged by WHO to approach their practice and research considering diverse human experience and intersectionality of various factors, including culture, gender, immigration status, and ethnicity, which may be related to lower physical and mental health²³⁻²⁵. In order to offer an acceptable and relevant health service to people of a certain cultural background, health promotion interventions could be developed from scratch with cultural sensitivity or a less resource-consuming way can, for example, be tailoring an already existing intervention for specific cultural groups²⁶. This process is defined as cultural adaptation²⁷. Cultural adaptation could be adopted on surface structure modifications (pairing materials and messages to apparent features of the target population such as language) or deep structure modifications (concerning intersecting effects of social, cultural, and historical variables on the target behavior)^{28,29}. Culturally adapted face-to-face interventions are shown to be effective in smoking cessation^{30,31}, health education, and healthy eating³². Moreover, IMI developed for ethnic minorities and underserved populations are also shown to be accepted and effective in the promotion of various health behaviors such as physical activity³³ and healthy eating¹⁶.

Previous reviews included IMI for ethnic minority and historically underserved populations in developed countries^{22,34-36}. However, none of them specifically examined culturally adapted

¹Department of Clinical Psychology and Psychotherapy, Institute of Psychology and Education, Ulm University, Lise-Meitner-Str. 16, D-89081 Ulm, Germany. ²Department of Rehabilitation Psychology and Psychotherapy, Institute of Psychology, University of Freiburg, Engelbergerstr. 41, D-79085 Freiburg, Germany. ✉email: sumeyye.balci@uni-ulm.de

IMI for health promotion. The number of studies exploring the development and dissemination of culturally adapted health promotion IMI is increasing and thus a systematic review and meta-analysis seems timely. Hence, this review aims to systematically identify culturally adapted IMI on health promotion and explore their effectiveness among populations that is different from the original intervention's target group.

RESULTS

All the predefined characteristics of included articles are presented in Table 1.

Study selection

We identified a total of 20,012 records. After screening titles and abstracts and full-texts, 13 studies were included in the quantitative analyses. The main characteristics of the studies are outlined in Table 1. During the full-text screening, 38 studies were excluded due to lack of cultural adaptation, 29 studies were concerning culturally sensitive interventions, 13 studies did not report relevant outcome data, eight studies were no original articles, four studies were not RCTs, five were excluded due to other reasons (i.e. novel interventions, study protocols, non-English full-text) and one due to a differing health promotion topic.

Study characteristics

The 13 RCTs included in this review comprise a total of $N = 10,747$ randomized participants, mostly adult populations ($N = 9710$). Trials were conducted between 2012 and 2020^{37–49}. The mean age of the participants varied from 14 to 57 (see Table 1). Primary studies focused on smokers ($k = 3$, $N = 2546$), individuals diagnosed with Type 2 diabetes ($k = 2$, $N = 171$), and HIV + individuals ($k = 1$, $N = 433$).

Four studies dealt with smoking cessation^{37,39,41,49}, two with both healthy eating and physical activity^{38,42}, three with physical activity only^{45–47}, two with healthy eating only^{40,43}, one with sexual health behavior⁴⁴, and one with alcohol consumption⁴⁸.

Four studies provided follow-up data on short-term effectiveness (one to five months follow-up)^{39,41,42,46}, three studies on long-term effectiveness (six to 12-months follow-up)^{40,45,47}, six studies provided follow-up data on both assessment points.

Seven studies were conducted in the USA^{38,39,43–45,47,48}, two in China^{37,42}, two in Brazil^{40,41}, one in Hong Kong⁴⁶, and one in Australia⁴⁹.

The cultural adaptation was based on a theory or a guideline in three studies^{38–40}. Eight studies based their cultural adaptation on a formative research/pilot study or expert review^{37,41,42,44,45,47–49}. Two studies did not provide information regarding the basis of cultural adaptation^{43,46}. In terms of alterations of the intervention content, four studies incorporated both surface and deep structure changes^{38,40,43,45}, while nine studies used surface structure changes only²⁸. Details of the culturally adapted and original interventions are presented in Table 2.

Risk of bias of in included studies

The risk of bias assessment of the included studies is presented in Figs. 1, 2. The interrater reliability suggested substantial agreement between the raters, $\kappa = 0.79$. Five studies were assessed to have a low risk of bias, five studies had some risk of bias and three studies were rated to have a high risk of bias. Four studies were assessed to have some risk of bias due to deviations from the intended intervention^{40,43,46,48}, three studies were assessed to have some risk of bias in the measurement of the outcome domain^{41,44,46}. Two studies^{37,38} were assessed to have some risk of bias due to missing outcome data, one study⁴³ had a high risk of bias in this domain. One study was assessed to have some risk of

bias arising from the randomization process⁴⁰. This study had an unequal number of clusters, which resulted in significant baseline differences in primary outcomes.

Effectiveness

Nine studies that provided data to calculate standardized mean difference effect sizes were pooled. Individual data points are presented in Supplementary Table 1. Four studies concerning smoking cessation did only provide dichotomous outcomes. We conducted separate meta-analyses for the different areas of health promotion (physical activity, smoking cessation), in the case of at least three studies reported on the same outcome. Since we included only nine studies in the analysis, we refrained from exploring the publication bias via a funnel plot⁵⁰. Following, we report on pooled effectiveness across health promotion domains.

The meta-analysis of six studies^{38,43–45,47,48} examining any long-term health promotion interventions revealed that culturally adapted IMI of health promotion were not superior to active control conditions in the long-term. In addition, four studies^{38,43,44,48} that provided short-term follow-up data were not superior to active control conditions in improving health behavior outcomes. Due to substantial heterogeneity, the results are reported descriptively, see Figs. 3, 4.

Comparisons with passive control groups could not be pooled given the low number of studies ($k = 4$), see Fig. 5. One study³⁹ reported only Odds ratios ([OR], 1.13; 95% CI 0.18 to 7.04).

Due to a small number of studies ($k = 9$), we did not perform the predefined subgroup and sensitivity analyses.

Effectiveness of culturally adapted IMI of physical activity

Five studies reported physical activity outcomes, four of which provided data via accelerometers/pedometers^{38,45,47} and one via a self-reported questionnaire⁴⁶. Pooling the three studies with active control conditions resulted in a small significant long-term effect favoring culturally adapted IMI ($N = 296$; $g = 0.48$; 95% CI 0.25 to 0.71; $I^2 = 41\%$; fixed effect), see Fig. 6.

Effectiveness of culturally adapted IMI of smoking cessation

Three RCTs ($N = 8,112$) reported smoking cessation outcomes measuring short-term abstinence at the end of the intervention versus active controls. The meta-analysis findings of these studies were not significant (Odds Ratio [OR], 1.75; 95% CI 0.51 to 6.05, $I^2 = 56\%$), see Fig. 7. The number of included studies was small ($k = 3$) and the effect size was mainly based on one large-scale study with $N = 8000$ participants³⁷.

Effectiveness of culturally adapted IMI of healthy eating, sexual health behavior, and alcohol consumption

Fewer than three studies reported on sexual health behavior⁴⁴, alcohol consumption⁴⁸, and healthy eating^{40,43}. Only one generic outcome, health-related quality of life, was reported in one trial⁴². These studies were not eligible for pooling due to the number of trials in the respective outcome but reported in Figs. 3–5.

DISCUSSION

To our knowledge, this is the first systematic review and meta-analysis that assessed the effectiveness of culturally adapted IMI on health promotion among populations which the intervention was originally not developed for. Our results suggested that culturally adapted IMI are not more effective in promoting health behaviors than active conditions in short- ($g = 0.10$) and long-term ($g = 0.20$), respectively. When regarding health promotion topics separately, health promotion of physical activity resulted in a small to moderate effect favoring culturally adapted IMI over active

Table 1. Characteristics of included articles.

1st author (year)	Country	Sample	Sample size E: experiment group/C: control group	Gender Female (%)	Mean age (SD)	Dropout rate at post-assessment (%)	Website vs. Mobile	Duration/ No. modules	Post randomization follow up in months	Comparison	Outcome	Outcome measures
Augustson et al. ³⁷	China	Adults smokers	E: 4000 C: 4000	3.6	–	73	SMS	6 weeks	1, 3 & 6 M	Active control group The Low-Frequency Text Contact (LFTC) received 1 text message a week, for the 6-week intervention period	Smoking cessation	Smoking status was based on past-7-day abstinence self-reported via text message
Bender et al. ³⁸	USA	Individuals diagnosed with Type 2 Diabetes	E: 22 C: 23	62	57.6 (9.8)	2.5	App/Social media	26 weeks	3 & 6 M	Active control group receives only Fitbit accelerometer and training for daily wear.	Physical activity	Step count via the Fitbit Zip. (accelerometer data)
Bowen et al. ³⁹	USA	Students (6th to 12th graders)	E: 64 C: 49	53	14.6	9	Website	6 weeks	1 M	Waitlist control	Smoking cessation	Smoking status based on "A Smoking Prevention Interactive Experience (ASPIRE)" Instrument
Brito Beck da Silva et al. ⁴⁰	Brazil	Students (7 th to 9 th grade)	E: 428 C: 467	46	14.49 (1.42)	30	Website	16 weeks	12 M	Waitlist control	Healthy eating	BMI
Cruvinel et al. ⁴¹	Brazil	Adult smoker post-discharge patients	E: 44 C: 22	45	47.7 (11.5)	10	Mobile/ SMS	2 weeks	1&3 M	Treatment as usual includes educational materials, brief intervention (BI), and access to NRT (adhesive patch and gum)	Smoking cessation	Smoking status of smokers (cigarettes a day) and self-reported 7-day point prevalence abstinence post-randomization.
Duan et al. ⁴²	China	University students	E: 270 C: 223	60	19.3 (1.07)	45	Website	8 weeks	2&3 M	Waitlist control	Physical activity & Quality of life	Chinese short version of the International Physical Activity Questionnaire (IPAQ-C) & Hong Kong version of the WHO's Quality of Life-BREF questionnaire
Fortmann et al. ⁴³	USA	Individuals diagnosed with Type 2 Diabetes	E: 63 C: 63	75	48.43 (9.8)	10	SMS	26 weeks	3&6 M	Treatment as usual (standard diabetes care provided by primary care providers at the clinic and group Diabetes self-management education- use of these services based on patient and physician's initiative)	Healthy eating	BMI
Kurth et al. ⁴⁴	USA	HIV+ individuals	E: 226 C: 207	55	47.8	8	Website	52 weeks	3, 6 & 9 M	Active control group (received computer-based audio-narrated risk assessment, which included questions about sexual risk behaviors, substance use, mental health, social support, partner status and disclosure, ART regimen and adherence in last 7 and 30 days, and side effects.)	Sexual health behavior	Sexual transmission risk behaviors (lack of condom use with either a main or another partner)

Table 1 continued

1st author (year)	Country	Sample	Sample size E: experiment group/C: control group	Gender Female (%)	Mean age (SD)	Dropout rate at post-assessment (%)	Website vs. Mobile	Duration/ No. modules	Post randomization follow up in months	Comparison	Outcome	Outcome measures
Larsen et al. ⁴⁵	USA	Adult male	E: 22 C: 24	0	43.04 (10.67)	6	SMS	24 weeks	6 M	Active control group (wellness control group received two SMS weekly throughout the study and publicly available print-based materials on health topics different from physical activity)	physical activity	Minutes/week of moderate to vigorous PA (MVPA) measured by accelerometers
Lau et al. ⁴⁶	Hong Kong	Students aged between 12-16 years old	E: 13 C: 16	49	13.7	not reported	SMS	4 weeks	1 M	No treatment	physical activity	Self-reported physical activity via PAQ-C (Physical activity questionnaire)
Marcus et al. ⁴⁷	USA	Inactive adult Latinas	E: 104 C: 101	100	39.20 (10.47)	not reported	Website	26 weeks	6 M	Active control group (wellness contact, receive access to a Spanish language website with information on health topics different from physical activity)	physical activity	Minutes/week via 7-day Physical Activity Recall and accelerometers.
Montag et al. ⁴⁸	USA	American Indian/ Alaska Native women	E: 113 C: 134	100	28.6	6	Website	20 min	1,3 & 6 M	Treatment as usual (get access to displayed educational brochures about health apart from FASD (fetal alcohol spectrum disorders) related information in the various waiting areas)	Alcohol consumption	Level of alcohol consumption (number of drinks per week)
Peiris et al. ⁴⁹	Australia	Current Aboriginal smokers (>16 years old)	E: 25 C: 24	78	42 (14)	6	Mobile App	53 weeks	1&6 M	Active control group (encouraged to use any other smoking cessation service or support and were offered Quitline and local ACCHS (Aboriginal Community Controlled Health Services) contact numbers)	Smoking cessation	Smoking status, self-reported abstinence

^aProvided three intervention groups versus a control group comparison, we used the intervention group which had the most exposure to the intervention as a comparator.

Table 2. Summary of culturally adapted and original IMI.

1st author (year)	Name original IMI adapted IMI	Language		Target group		Ethnicity		Health promotion		Cultural adaptation theory	Cultural adaptation components
		original IMI	adapted IMI	original IMI	adapted IMI	original IMI	adapted IMI	original IMI	adapted IMI		
Augustson et al. ³⁷		English	English	General population	US American	Smoking cessation	Smoking cessation	Expert review, focus groups	Language, context adaptation		
Bender et al. ³⁸	Change to Quit China Diabetes Prevention Program (DPP)	Chinese	English	General population Type 2 Diabetes patients	Chinese American	Smoking cessation	Healthy eating/ physical activity	Bender & Clark (2011)'s theory ¹⁰⁷	Content (Filipino food photos), delivery (involvement of family members to the office visits) language		
Bowen et al. ³⁹	PiAm Go4Health SmokingZine	English	English	Type 2 Diabetes patients General population (adolescent)	Filipino Canadian	Healthy eating/ physical activity	Smoking cessation	Based on a guideline from Wisdom2Action	Images, context		
Brito Beck da Silva et al. ⁴⁰	- StayingFit	English	English	General population (adolescents) General population	American Indian US American	Smoking cessation	Smoking cessation	Based on Barrera et al (2013) ¹⁰⁸ and Castro et al (2015) ¹⁰⁹	Language, cultural standards, meanings, and values added		
Cruvinel et al. ⁴¹	- TXT StayingFit Brazil	Portuguese	Portuguese	General population (adolescents) Hospitalized smokers	Brazilian US American	Smoking cessation	Smoking cessation	Formative research	Language, information from the Brazilian smoking cessation treatment guideline		
Duan et al. ⁴²	-	-	-	General population	US, Germany and Netherlands	Healthy eating/ physical activity	Smoking cessation	Formative research	Language, content		
Fortmann et al. ⁴³	- Staged Diabetes Management (SDM) & Dulce Project Dulce Digital	Chinese	English	General population (students) General population	Chinese US American	Healthy eating/ physical activity	Healthy eating/ diabetes management	Based on face-to-face intervention project Dulce ¹¹⁰	Language, cultural beliefs that interfere with optimum self-management, shortened content, motivational messages		
Kurth et al. ⁴⁴	- CARE + CARE + Spanish	English and Spanish	English	Type 2 Diabetes patients HIV + patients	Hispanic US American	Healthy eating	Sexual health behavior	The local expert advisory panel, usability testing	Content (Language), expert suggestions		
Larsen et al. ⁴⁵	- Seamos Saludables Activo	Spanish	Spanish	General population General population (men)	US American Latino	Sexual health behavior	physical activity	Formative research and pilot (qualitative interviews)	Language adaptation, the content of the SMS, and printed materials		
Lau et al. ⁴⁶	-	English	English, Dutch, Turkish	General population General population	US American/Canadian Hong Kong Chinese	physical activity	physical activity	NA	Language, content (colloquial dialogue for adolescents)		
Marcus et al. ⁴⁷	- Pasos Hacia la Salud	English	English	General population General population (women)	US American Latina	physical activity	physical activity	Focus groups	Cultural and linguistic adaptation, culturally adapted content and support specifically for Latinas, flexible scheduling for assessment meetings, reimbursement for travel and childcare		

Table 2 continued

1st author (year)	Name		Language		Target group		Ethnicity		Health promotion		Cultural adaptation theory	Cultural adaptation components
	original IMI	adapted IMI	original IMI	adapted IMI	original IMI	adapted IMI	original IMI	adapted IMI	original IMI	adapted IMI		
Montag (2015)	e-CHUG		English		General population		US American		Alcohol consumption		Focus groups	Content (pictures, logo, color of the layout, example characters, myths) - added video (verbal tradition)- language (not a translation but wording and simplifying)
	eCHECKUP TO GO		English		General population (women)		American Indian/ Alaska Native (AIAN)		Alcohol consumption			
Peiris et al. ⁴⁹	QuitTxt		English		General population		Australian		Smoking cessation		Formative research with the expert user group	Adaptation of the content and tone of the messages based on the attitudes of the target group towards smoking
	Can't Even Quit'		English		General population		Australian/Aboriginal/ Citizen of Torres Strait Island		Smoking cessation			

control conditions in the long-term ($g = 0.48$). This is in line with a previous umbrella review of health promotion IMI for minority and historically underserved populations, which, however, not exclusively included culturally adapted IMI²². Similarly, another meta-analysis of IMI concerning physical activity did highlight the superiority of IMI over a control group or no-treatment condition, without a specific focus on culturally adapted interventions¹⁴. No other significant effect was revealed for the other addressed health promotion topics.

Subgroup analyses aiming to detangle the substantial between trial heterogeneity were not feasible. Heterogeneity among the included studies was moderate to substantial, I^2 ranged from 0.36 to 0.66. Prior research points to differential effects on culturally adapted health promotion interventions in terms of different populations (age and ethnicity⁵¹), different intervention features (professional vs. non-professional provider)³², intervention duration and follow-up times in culturally adapted face-to-face interventions⁵², interventions focusing on general population groups⁵³, different methodological decisions (RCT methodology use, different control groups, e.g. tailored website, no-treatment controls⁵³) and cultural adaptation contents (inclusion of social support and/or family members⁵¹, integrating cultural beliefs and values⁵⁴). There might be further possible explanations for between-study heterogeneity, and future research needs to provide a better understanding of the impact such factors have on the effectiveness of culturally adapted IMI for health promotion. The present findings are inconsistent with previous meta-analyses on IMI in several ways. Among western populations, these meta-analyses yielded positive effects favoring IMI compared to a waitlist and/or active controls (e.g. other internet-based or face-to-face interventions): for smoking cessation and abstinence outcome^{10,55}, which is maintained at 12-month follow-up⁵⁶ and higher effects achieved with the use of tailored messages^{10,13}; healthy eating^{57–59} and weight loss⁶⁰; sexual health behavior promotion⁶¹ and regarding HIV prevention and condom use⁶². Another meta-analysis found significant positive effects of tailored (based on personal relevance) web-based interventions on health behaviors compared to non-tailored web-based interventions⁵³ and a different meta-analysis of SMS-based interventions on various health behavior outcomes suggested that targeted and tailored (based on demographic and psychosocial factors) SMS yielded larger effect sizes, especially for physical activity interventions ($g = 0.51$), which yielded a similar effect size to our results ($g = 0.48$)¹⁰. Most surprisingly in this context is our null-finding regarding the effectiveness of culturally adapted IMI compared to waitlist control conditions, which are known to provide a rather upper benchmark of the benefit of interventions⁶³, usually associated with significantly larger between-group effect sizes in IMI for health promotion as well^{10,13,64}. Although some of the meta-analyses mentioned above concern tailored intervention contents, none of the above-mentioned meta-analyses were specifically examining culturally adapted IMI. Therefore, our results cannot be easily compared with prior meta-analyses. However, if culturally adapted IMI for health promotion are not effective at all, even when compared to waitlist controls, we might need to challenge the idea of providing culturally adapted IMI to populations for which the intervention was originally not developed for at large and examine whether IMI developed with cultural sensitivity are effective in the same target groups. Hence, explanations for this surprisingly limited effectiveness seem warranted.

One possible explanation of our results might be related to the quality of cultural adaptation of the interventions. The cultural adaptation processes were rarely well defined in the included studies. Therefore, it was not clear whether aspects of cultural adaptation were appropriate. In addition, the high dropout among included studies could be an indicator of cultural adaptation not




Fig. 1 Risk of bias summary. Reviewers' judgments about each risk of bias item for each included study.

working as intended by the researcher. Moreover, only three studies based their cultural adaptation process on a theory, which might contribute to its quality. A tested theory of cultural adaptation of IMI is missing. However, there are guidelines developed for culturally adapting face-to-face interventions^{28,65} and researchers could implement these guidelines when adapting an IMI⁶⁶. A recent taxonomy of cultural adaptation of IMI for mental disorders serves as a basis for future cultural adaptations of IMI⁶⁷. Adopting a theoretical basis in intervention development is suggested to result in higher effects, as was shown in a meta-analysis¹⁴. However, this could not be shown in our results: only one out of the three IMI that utilized a theory of cultural adaptation resulted in an improvement in physical activity outcome³⁸. In the future, culturally adapted intervention studies should consider supporting cultural adaptation with an established theory and report the adaptation process in more detail to lead prospective cultural adaptations and replications. In the process of cultural adaptation, some of the included studies sought expert reviews, focus group feedback, and conducted a pilot study, and at least altered one aspect of the intervention. However, the majority of the changes were regarded as taking place only at the surface structure²⁸, which might be one reason for the limited impact of culturally adapted IMI shown in the present review. Surface structure changes aim at improving feasibility while deep structure changes target program's effect for the participants²⁸. Implementing deep structure changes involves core cultural values of a certain population, such as beliefs towards a health issue, or performing a behavior as a member of gender identity. A meta-analysis of face-to-face culturally adapted health interventions showed that incorporating family members and religious values, which are considered as deep structure changes, in the intervention was related to improvements³¹. Future research should consider exploring surface versus deep structure alterations on the effectiveness of IMI e.g. in the framework of dismantling and additive clinical trial designs aiming at detangling active components and mechanisms of change of the respective interventions^{68–71}.

Moreover, culturally adapting interventions is not free from criticism. It should be taken into account that the majority of the interventions developed in the fields of psychology and behavioral medicine are for a rather homogeneous group (white, educated, middle to high socioeconomic status) but not representative for the majority^{19,72}. In the cultural adaptation process, the same intervention is often altered to cater to the needs of a different group of people that are non-white, occasionally less educated, and/or bilingual. This process could be seen as a form of assimilation for the target group because even the topic of the intervention might also be representative of western, white, educated humans¹. Therefore, in order to avoid these issues, the first step of cultural adaptation might include approaching the problem and defining it with the cultural sensitivity of its target group⁷³. In addition, people of a certain

cultural background are not homogenous within themselves, each member's experience is affected by intersecting factors⁷⁴. Therefore, it might be more complicated than often expressed to adapt an intervention for a cultural group⁷⁵. One possible solution might be to invest in adapting interventions to cultural specifics of the users, e.g. based on user needs assessments⁷⁶ or community leaders' input⁷⁷. Another solution, especially for migrant/immigrant populations could be developing interventions based on the target groups' acculturation levels, i.e. a process on a spectrum of either orientation to the host culture or maintaining the native culture^{1,76}. It seems also worthwhile to pay attention to intersecting factors that might influence a member of a cultural group, namely gender and literacy. However, we first need to establish whether to culturally adapt health promotion IMI at all. The present findings at least suggest—except for physical activity IMI—a non-favorable cost-benefit ratio, a result that still is in need of stronger evidence.

Another topic of relevance to our findings is the reach, uptake, and intervention adherence of culturally adapted IMI. Even if such IMI would be effective, they still need to be used in order to exploit their full potential. Although internet technologies are widely used globally⁷⁸, there are still barriers to utilizing these technologies, which cause inequalities in accessing the internet and mobile technologies and comprehending health information^{16,79,80}. Moreover, pure mobile-based interventions are seemingly less effective than internet-based or combined interventions^{81–83}, which might also affect for instance minority populations, where they are more likely to use a smartphone to access the internet than non-minority populations where multiple device (e.g. tablets, desktops) ownership is common⁸⁴. To increase adherence, multimodal content and guidance (direct contact with the provider)⁸⁵ might be useful via diminishing issues of health literacy, motivational and volitional aspects, and the digital divide^{71,86,87}. These aspects might be particularly important in people of certain cultural backgrounds living in a high-income country, people living in low-income countries, and/or vulnerable populations, such as immigrants, given the limited representation of many of these populations in the health research, and high rates of drop-out^{86,88–91}.

This review has some limitations. First, we included interventions concerning only five prominent areas of health promotion thus results might not be generalizable to other health promotion domains. Second, the definition of cultural adaptation varies, and our operationalization of culturally adapted interventions resulted in the exclusion of studies that investigated IMI that were developed newly in a culturally sensitive way. Comparison of culturally adapted vs. culturally sensitive interventions is an interesting further research topic. Moreover, comparing culturally adapted versus culturally sensitive IMI might present insights into whether it is worthwhile to develop a novel IMI for a group or adapt an already existing one. Third, we were able to pool data from only 13 studies, which further limit the generalizability of our




Fig. 2 Risk of bias graph. Reviewers' judgments about each risk of bias item presented as percentages across all included studies.

results. Due to this limited evidence base, analyses were restricted to the main research questions while subgroup analyses were not feasible yet. Future updates might allow for exploring the between-study heterogeneity highlighted in the present review,

while the findings reported here could guide researchers in what to examine next. In this context, we suggest adding generic outcomes such as health-related quality of life, daily functioning, or self-efficacy to domain/disease-specific outcomes to allow for




Fig. 3 Summary of culturally adapted IMI of health promotion vs. active controls in the long-term. Due to substantial heterogeneity among the culturally adapted IMI of health promotion vs. active controls in long-term meta-analytical pooling did not perform.




Fig. 4 Summary of culturally adapted IMI of health promotion vs. active controls in the short-term. A summary plot of effect sizes of four studies of culturally adapted IMI of health promotion vs. active controls in short-term are presented.




Fig. 5 Summary of culturally adapted IMI of health promotion vs. passive controls. Due to few numbers of studies (two studies reported data in the long-term, two in the short-term, while one study reported dichotomous outcome) comparing culturally adapted IMI to a passive control group, meta-analytic pooling did not perform.

cross-trial cross health promotion domain comparisons in the future. Fourth, although the studies included in this meta-analysis were culturally adapted, the adaptation process was rarely well defined. Fifth, only three out of 13 studies used a theory to adapt the intervention. And last, none of the studies were comparing culturally adapted IMI to non-adapted IMI. This creates a difficulty to draw any firm conclusions about the differential effectiveness of culturally adapted interventions. Despite these limitations, this meta-analysis had some strengths. To our knowledge, this is the




Fig. 6 Fixed effects meta-analysis of culturally adapted IMI for physical activity vs. active control conditions. Forest plot presenting fixed effects meta-analysis of culturally adapted IMI for physical activity vs. active controls.

first meta-analysis concerning culturally adapted health promotion IMI. Moreover, this a-prior registered systematic review included mostly scientifically sound RCTs from a broad sample and health behavior topics. A summary of recommendations for future research is represented in Box 1.

CONCLUSION

Based on the present findings, culturally adapted IMI might not be superior compared to control conditions in the short- and long-term, except for physical activity. Although they might exhibit a more attractive health offer to their target group, their usefulness is questionable or at least need further examination. Thereby, it might be worthwhile to take into consideration intersecting aspects of experiences of people of certain cultural groups regarding health behaviors to assure acceptability and effectiveness when designing interventions and contribute to diminishing health inequalities.

METHODS

Protocol and registration

This systematic review and meta-analysis has been registered at PROSPERO (Registration number: CRD 42020152939) and follows the format of the PRISMA guideline⁹². Review protocol⁹³ described the aim, methodology, and data analysis plan in advance. Changes to study protocol are listed in the supplementary notes.

Eligibility criteria

Studies were included if they (1) were RCTs, (2) had no treatment, treatment as usual (TAU), placebo, waitlist, or active control conditions, (3) were delivered via the internet, (4) were culturally adapted for a population that differed from the original intervention's target group, (5) examined a health promotion intervention on healthy eating, physical activity, alcohol consumption, sexual health behavior and/or smoking cessation (6) reported one of the respective health promotion-specific outcomes: body mass index (BMI), time spent exercising, change in condom use, level of smoking, level of alcohol consumption, or one of the following generic outcomes: health-related quality of life and self-efficacy.

Information sources and study selection

The initial search was conducted in the following databases on 26. August 2019: Cochrane Central Register of Controlled Trials (CENTRAL), EbscoHost/MEDLINE, Ovid/Embase, EbscoHost/PsychINFO, and Web of Science. A combination of keywords (including MeSH terms) indicating culturally adapted IMI for health promotion has been used. The search terms are published in this review's protocol⁹³ and cover comprehensively both the topic of the present review as well as one of a parallel systematic




Fig. 7 Summary of culturally adapted IMI for smoking cessation vs. active controls in short-term. Three studies reported smoking cessation outcomes measuring short-term abstinence at the end of the intervention vs. active controls are presented on the forest plot.

Box 1 Recommendations for future research

1. Improve evidence on whether culturally adapted IMI for health promotion are indeed not effective at all, even when compared to passive control conditions.
2. Provide a better understanding of the impact of population and intervention characteristics on the differential effects of culturally adapted IMI for health promotion.
If culturally adapted IMI are effective at least compared to passive control conditions (see 1.) and/or at least with regard to some subgroups (see 2.):
3. Improve evidence on whether culturally adapted IMI are effective compared to active control conditions.
4. Particularly compare culturally adapted IMI with the respective non-adapted versions or simple language translations of the IMI.
Ultimately, the substantial effort necessary for adapting IMI culturally might only be justified in case of clinically significant superiority of the culturally adapted versions compared to active controls. In case this is given (see 3./4.)- at least with regard to some subgroups (see 2.):
5. Examine active components and mechanisms of change of these effective culturally adapted IMI. Particularly provide a better understanding of the impact of surface and deep structure changes on intervention adherence and effectiveness.
6. Examine ways of improving reach, uptake, engagement, and intervention adherence for the effective culturally adapted IMI.
7. Develop evidence-based recommendations and guidelines for adapting IMI culturally delineated from the effective culturally adapted IMI.

review on culturally adapted IMI for mental health conditions^{94,67}. There were no restrictions on the publication date. A search update was conducted on 15 October 2020.

All search results were merged into Covidence⁹⁵ and duplicates were automatically removed. Two reviewers screened titles and abstracts of the identified articles against the inclusion criteria and selected potentially relevant articles for the full-text screening. Full-text screening has been performed by two reviewers independently, disagreements have been solved by consensus or a third reviewer where needed. The study selection is illustrated in the PRISMA flow diagram (see Fig. 8).

Data extraction

Data extraction was conducted by two independent reviewers and then extracted data was then checked by a third reviewer. The following data were extracted from the included studies: publication details, study participants (demographics and cultural background, baseline characteristics), study design, study setting, characteristics of the original and culturally adapted intervention, health behavior-specific and generic outcome measures, information regarding cultural adaptation (content, utilization of theoretical or evidence-based components). Behavioral outcomes are defined as: physical activity measured via physical activity minutes per week with accelerometers or self-report questionnaires; healthy eating measured via BMI; alcohol consumption assessed via the level of alcohol

consumption; smoking cessation assessed via the level of smoking or the abstinence percentage; sexual health behavior assessed with condom use. Generic outcomes were defined as health-related quality of life and self-efficacy, assessed by means of validated self-report questionnaires. Control conditions were categorized into active (placebo, other health promotion interventions & TAU) and passive controls (waitlist & No treatment). When related information could not be extracted, corresponding authors of the articles were contacted to obtain information. The extracted data was tabulated.

Risk of bias

Two independent reviewers performed quality assessments with Cochrane Collaboration's Risk of Bias Tool 2.0⁹⁶. The RoB tool 2.0 has five domains including bias arising from the randomization process, bias due to deviations from intended interventions, bias due to missing outcome data, bias in the measurement of the outcome, bias in the selection of the reported result. A third reviewer solved disagreements following a discussion between the reviewers. The Kappa statistic was used to calculate interrater reliability⁹⁷.

Meta-analysis

For each study, a standardized mean difference (SMD) and 95% confidence intervals (CI) were calculated with mean scores of intervention and control groups. In order to decrease the bias of small samples, Hedges' g was calculated⁹⁸. Effect sizes were recoded when higher scores of an outcome assessment indicated worsening results (e.g. BMI and level of alcohol consumption). For continuous outcomes, Hedges' g and 95% CIs were reported; for dichotomous outcomes, odds ratios and CIs were reported. Random effects model was chosen for analyses due to an expected diversity among IMI of health promotion, sample size, and duration of intervention among studies⁹⁹. Data were pooled to calculate a standardized mean effect size for each outcome and a forest plot with 95% CIs, in the case of at least three studies reporting the respective outcome. Otherwise, results were presented descriptively. Sensitivity analysis was planned to assess the impact of studies with a high risk of bias. Analyses were performed in R package meta and metafor, and Review Manager 5¹⁰⁰⁻¹⁰³. Continuous effect sizes were categorized along with Cohen's rule of thumb with 0.20 considered a small effect, 0.50 medium effect, and 0.80 large effects¹⁰⁴. In order to assess publication bias, we planned to conduct funnel plots.

Statistical heterogeneity among studies was analyzed with the I^2 statistics^{97,105}. Statistical heterogeneity refers to the variability among effect sizes in a meta-analysis¹⁰⁶. However, the veracity of measures of heterogeneity is arguable; therefore, their interpretation should be made with caution⁹⁷. Statistical heterogeneity was calculated with the I^2 test for each outcome domain. Heterogeneity $I^2 \geq 60\%$ was regarded as substantial heterogeneity, in which case no pooled effect sizes are reported. Subgroup analyses were planned to explore possible sources of




Fig. 8 Prisma Flow chart⁹². Study identification, selection, and inclusion represented on the diagram. An asterisk symbol represents a parallel review conducted regarding the culturally adapted internet- and mobile-based interventions concerning mental health.

heterogeneity on population, duration of the intervention, delivery of the intervention, and presence of guidance.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary linked to this article.

DATA AVAILABILITY

Data supporting the findings of this study are available within the respective articles cited in this review and from the corresponding author on reasonable request.

CODE AVAILABILITY

Meta-analyses were conducted using the R Studio packages Meta¹⁰⁰ and Metafor¹⁰³, and Review Manager^{101,102}. The code is publicly available in the following github repository: <https://github.com/MathiasHarrer/Doing-Meta-Analysis-in-R>.

Received: 18 August 2021; Accepted: 21 January 2022;

Published online: 23 March 2022

REFERENCES

1. Liu, J. J. et al. Adapting health promotion interventions to meet the needs of ethnic minority groups: Mixed-methods evidence synthesis. *Health Technology Assessment* <https://doi.org/10.3310/hta16440> (2012).
2. Cerf, M. E. Healthy lifestyles and noncommunicable diseases: nutrition, the life-course, and health promotion. *Lifestyle Med.* **2**, 1–12 (2021).
3. Preston, S. H., Stokes, A., Mehta, N. K. & Cao, B. Projecting the effect of changes in smoking and obesity on future life expectancy in the United States. *Demography* **51**, 27–49 (2014).
4. World Health Organization. *Sexual Health and its Linkages to Reproductive Health: an Operational Approach* 1–12 (World Health Organization, 2017).
5. Di Cesare, M. et al. Inequalities in non-communicable diseases and effective responses. *Lancet* **381**, 585–597 (2013).
6. World Health Organization. *Global Status Report On Noncommunicable Diseases 2014* (World Health Organization, 2014).
7. World Health Organization. *2008-2013 Action Plan for the Global Strategy for the Prevention and Control of Noncommunicable Diseases* (World Health Organization, 2013).
8. Friel, S. & Marmot, M. G. Action on the Social determinants of health and health inequities goes global. *Annu. Rev. Public Health.* <https://doi.org/10.1146/annurev-publhealth-031210-101220> (2011).
9. Muñoz, R. F. Using evidence-based Internet interventions to reduce health disparities worldwide. *J. Med. Internet Res.* **12**, 1–10 (2010).

10. Head, K. J., Noar, S. M., Iannarino, N. T. & Grant Harrington, N. Efficacy of text messaging-based interventions for health promotion: a meta-analysis. *Soc. Sci. Med.* **97**, 41–48 (2013).
11. Fiedler, J., Eckert, T., Wunsch, K. & Woll, A. Key facets to build up eHealth and mHealth interventions to enhance physical activity, sedentary behavior and nutrition in healthy subjects—an umbrella review. *BMC Public Health* **20**, 1–21 (2020).
12. Müller, A. M. et al. Physical activity, sedentary behavior, and diet-related ehealth and mhealth research: Bibliometric analysis. *J. Med. Internet Res.* **20**, e122 (2018).
13. Do, H. P. et al. Which eHealth interventions are most effective for smoking cessation? A systematic review. *Patient Prefer. Adherence* **12**, 2065–2084 (2018).
14. Webb, T. L., Joseph, J., Yardley, L. & Michie, S. Using the Internet to promote health behavior change: a systematic review and meta-analysis of the impact of the theoretical basis, use of behavior change techniques, and mode of delivery on efficacy. *J. Med. Internet Res.* **12**, 1–18 (2010).
15. Muñoz, R. F. et al. Massive open online interventions: a novel model for delivering behavioral health services worldwide. *Clin. Psychol. Sci.* **4**, 194–205 (2016).
16. Vandelanotte, C. et al. Past, present, and future of ehealth and mhealth research to improve physical activity and dietary behaviors. *J. Nutr. Educ. Behav.* **48**, 219–228.e1 (2016).
17. Jiang, X., Ming, W. K. & You, J. H. S. The cost-effectiveness of digital health interventions on the management of cardiovascular diseases: systematic review. *J. Med. Internet Res.* **21**, 1–11 (2019).
18. Elbert, N. J. et al. Effectiveness and cost-effectiveness of ehealth interventions in somatic diseases: a systematic review of systematic reviews and meta-analyses. *J. Med. Internet Res.* **16**, 1–23 (2014).
19. Henrich, J., Heine, S. J. & Norenzayan, A. Beyond WEIRD: towards a broad-based behavioral science. *Behav. Brain Sci.* **33**, 111–135 (2010).
20. Lin, J., Faust, B., Ebert, D. D., Krämer, L. & Baumeister, H. A web-based acceptance-facilitating intervention for identifying patients' acceptance, uptake, and adherence of internet- and mobile-based pain interventions: randomized controlled trial. *J. Med. Internet Res.* **20**, e244 (2018).
21. Baumeister, H. et al. Impact of an acceptance facilitating intervention on diabetes patients' acceptance of Internet-based interventions for depression: a randomized controlled trial. *Diabetes Res. Clin. Pract.* **105**, 30–39 (2014).
22. Armaou, M., Araviaki, E. & Musikanski, L. eHealth and mHealth interventions for ethnic minority and historically underserved populations in developed countries: an umbrella review. *Int. J. Community Well-Being* **3**, 193–221 (2020).
23. Clauss-Ehlers, C. S., Chiriboga, D. A., Hunter, S. J., Roysircar, G. & Tummala-Narra, P. APA multicultural guidelines executive summary: ecological approach to context, identity, and intersectionality. *Am. Psychol.* **74**, 232–244 (2019).
24. Lewis, J. A., Williams, M. G., Peppers, E. J. & Gadson, C. A. Applying intersectionality to explore the relations between gendered racism and health among black women. *J. Couns. Psychol.* **64**, 475–486 (2017).
25. World Health Organization. Promoting health in the SDGs. *Report on 9th Global Conference on Health Promotion* 1–37 (World Health Organization, 2016).
26. Barrera, M., Castro, F. G. & Steiker, L. K. H. A critical analysis of approaches to the development of preventive interventions for subcultural groups. *Am. J. Community Psychol.* **48**, 439–454 (2011).
27. Bernal, G., Jiménez-Chafey, M. I., & Domenech Rodríguez, M. M. & Domenech Rodríguez Cultural adaptation of treatments: a resource for considering culture in evidence-based practice. *Prof. Psychol. Res. Pract.* **40**, 361–368 (2009).
28. Resnicow, K., Soler, R., Braithwaite, R. L., Ahluwalia, J. S. & Butler, J. Cultural sensitivity in substance use prevention. *J. Community Psychol.* **28**, 271–290 (2000).
29. Cardemil, E. Cultural adaptations to empirically supported treatments: a research agenda. *Sci. Rev. Ment. Heal. Pract.* **7**, 8–21 (2010).
30. Liu, J. J. et al. Smoking cessation interventions for ethnic minority groups—a systematic review of adapted interventions. *Prev. Med. (Balt.)* **57**, 765–775 (2013).
31. Nierkens, V. et al. Effectiveness of cultural adaptations of interventions aimed at smoking cessation, diet, and/or physical activity in ethnic minorities. a systematic review. *PLoS ONE* <https://doi.org/10.1371/journal.pone.0073373> (2013).
32. Lambert, S. et al. The effect of culturally-adapted health education interventions among culturally and linguistically diverse (CALD) patients with a chronic illness: a meta-analysis and descriptive systematic review. *Patient Educ. Couns.* <https://doi.org/10.1016/j.pec.2021.01.023> (2021).
33. Staffileno, B. A. B. A., Tangney, C. C. C. & Fogg, L. Favorable outcomes using an eHealth approach to promote physical activity and nutrition among young African American women. *J. Cardiovasc. Nurs.* **33**, 62–71 (2018).
34. Anderson-Lewis, C., Darville, G., Mercado, R. E., Howell, S. & Di Maggio, S. mHealth technology use and implications in historically underserved and minority populations in the united states: Systematic literature review. *JMIR mHealth uHealth* **6**, e128 (2018).
35. Montague, E. & Perchonok, J. Health and wellness technology use by historically underserved health consumers: Systematic review. *J. Med. Internet Res.* **14**, e78 (2012).
36. Bennett, G. G. et al. Electronic health (eHealth) interventions for weight management among racial/ethnic minority adults: a systematic review. *Obes. Rev.* **15**, 146–158 (2014).
37. Augustson, E. et al. Text to quit China: an mHealth smoking cessation. *Trial Am. J. Health Promot.* **31**, 217–225 (2017).
38. Bender, M. S., Cooper, B. A., Park, L. G., Padash, S. & Arai, S. A feasible and efficacious mobile-phone based lifestyle intervention for Filipino Americans with type 2 diabetes: randomized controlled trial. *JMIR Diabetes* **2**, e30 (2017).
39. Bowen, D. J., Henderson, P. N., Harvill, J. & Buchwald, D. Short-term effects of a smoking prevention website in American Indian youth. *J. Med. Internet Res.* **14**, e81 (2012).
40. Brito Beck Da Silva, K. et al. Evaluation of the computer-based intervention program stayingfit Brazil to promote healthy eating habits: The results from a school cluster-randomized controlled trial. *Int. J. Environ. Res. Public Health* **16**, 1–18 (2019).
41. Cruvinel, E., Richter, K. P., Colugnati, F. & Ronzani, T. M. An experimental feasibility study of a hybrid telephone counseling/text messaging intervention for post-discharge cessation support among hospitalized smokers in Brazil. *Nicotine Tob. Res.* **21**, 1700–1705 (2019).
42. Duan, Y. P., Wienert, J., Hu, C., Si, G. Y. & Lippke, S. Web-based intervention for physical activity and fruit and vegetable intake among Chinese university students: a randomized controlled trial. *J. Med. Internet Res.* **19**, 1–15 (2017).
43. Fortmann, A. L. et al. Dulce digital: An mHealth SMS based intervention improves glycemic control in hispanics with type 2 diabetes. *Diabetes Care* **40**, 1349–1355 (2017).
44. Kurth, A. E. et al. Linguistic and cultural adaptation of a computer-based counseling program (CARE+ Spanish) to support HIV treatment adherence and risk reduction for people living with HIV/AIDS: a randomized controlled trial. *J. Med. Internet Res.* **18**, e195 (2016).
45. Larsen, B. A. et al. Randomized trial of a physical activity intervention for Latino Men: Activo. *Am. J. Prev. Med.* **59**, 219–227 (2020).
46. Lau, P. W. C., Pitkethly, A. J., Leung, B. W. C., Lau, E. Y. & Wang, J. J. The intervention effect of SMS delivery on chinese adolescent's physical activity. *Int. J. Environ. Res. Public Health* **16**, 1–12 (2019).
47. Marcus, B. H. et al. Pasos Hacia La Salud: a randomized controlled trial of an internet-delivered physical activity intervention for Latinas. *Int. J. Behav. Nutr. Phys. Act.* **13**, 62 (2016).
48. Montag, A. C. et al. Preventing alcohol-exposed pregnancy among an american indian/alaska native population: effect of a screening, brief intervention, and referral to treatment intervention. *Alcohol. Clin. Exp. Res.* **39**, 126–135 (2015).
49. Peiris, D. et al. A smartphone app to assist smoking cessation among aboriginal australians: findings from a pilot randomized controlled trial. *JMIR mHealth uHealth* **7**, e12745 (2019).
50. Rothstein, H. R., Sutton, A. J. & Borenstein, M. *Publication Bias in Meta-Analyses: Prevention, Assessment and Adjustments* (John Wiley & Sons, 2005).
51. Rathod, S. et al. The current status of culturally adapted mental health interventions: a practice-focused review of meta-analyses. *Neuropsychiatr. Dis. Treat.* **14**, 165–178 (2018).
52. Harper Shehadeh, M., Maercker, A., Heim, E., Chowdhary, N. & Albanese, E. Cultural adaptation of minimally guided interventions for common mental disorders: a systematic review and meta-analysis. *JMIR Ment. Heal* **3**, e44 (2016).
53. Lustria, M. L. A. et al. A meta-analysis of web-delivered tailored health behavior change interventions. *J. Health Commun.* <https://doi.org/10.1080/10810730.2013.768727> (2013).
54. Van Loon, A., Van Schaik, A., Dekker, J. & Beekman, A. Bridging the gap for ethnic minority adult outpatients with depression and anxiety disorders by culturally adapted treatments. *J. Affect. Disord.* **147**, 9–16 (2013).
55. Whittaker, R. et al. Mobile phone text messaging and app-based interventions for smoking cessation. *Cochrane Database Syst. Rev.* <https://doi.org/10.1002/14651858.CD006611.pub5> (2019).
56. Myung, S.-K., McDonnell, D. D., Kazinets, G., Seo, H. G. & Moskowitz, J. M. Effects of web- and computer-based smoking cessation programs. *Arch. Intern. Med.* <https://doi.org/10.1001/archinternmed.2009.109> (2009).
57. Joiner, K. L., Nam, S. & Whittemore, R. Lifestyle interventions based on the diabetes prevention program delivered via eHealth: a systematic review and meta-analysis. *Prev. Med. (Balt.)* **100**, 194–207 (2017).
58. Rodriguez Rocha, N. P. & Kim, H. eHealth interventions for fruit and vegetable intake: a meta-analysis of effectiveness. *Heal. Educ. Behav.* **46**, 947–959 (2019).
59. Kelly, J. T., Reidlinger, D. P., Hoffmann, T. C. & Campbell, K. L. Telehealth methods to deliver dietary interventions in adults with chronic disease: a systematic review and meta-analysis. *Am. J. Clin. Nutr.* **104**, 1693–1702 (2016).
60. Hutchesson, M. J. et al. eHealth interventions for the prevention and treatment of overweight and obesity in adults: A systematic review with meta-analysis. *Obes. Rev.* **16**, 376–392 (2015).

61. Bailey, J. V. et al. Interactive computer-based interventions for sexual health promotion. *Cochrane Database Syst. Rev.* <https://doi.org/10.1002/14651858.cd006483.pub2> (2010).
62. Noar, S. M., Black, H. G. & Pierce, L. B. Efficacy of computer technology-based HIV prevention interventions: a meta-analysis. *Aids* **23**, 107–115 (2009).
63. Cunningham, J. A., Kypri, K. & McCambridge, J. Exploratory randomized controlled trial evaluating the impact of a waiting list control design. *BMC Med. Res. Methodol.* **13**, 150 (2013).
64. Graham, A. et al. Systematic review and meta-analysis of Internet interventions for smoking cessation among adults. *Subst. Abuse Rehabil.* **55**. <https://doi.org/10.2147/sar.s101660> (2016).
65. Bernal, G. & Sáez-Santiago, E. Culturally centered psychosocial interventions. *J. Community Psychol.* **34**, 121–132 (2006).
66. Salamanca-Sanabria, A., Richards, D. & Timulak, L. Adapting an internet-delivered intervention for depression for a Colombian college student population: an illustration of an integrative empirical approach. *Internet Inter.* **15**, 76–86 (2019).
67. Spanhel, K. et al. Cultural adaptation of internet- and mobile-based interventions for mental disorders: a systematic review. *npj Digit. Med.* <https://doi.org/10.1038/s41746-021-00498-1> (2021).
68. Domhardt, M. et al. Therapeutic processes in digital interventions for anxiety: a systematic review and meta-analytic structural equation modeling of randomized controlled trials. *Clin. Psychol. Rev.* **90**, 102084 (2021).
69. Domhardt, M., Cuijpers, P., Ebert, D. D. & Baumeister, H. More light? opportunities and pitfalls in digitalized psychotherapy process research. *Front. Psychol.* **12**, 1–5 (2021).
70. Kazdin, A. E. Mediators and mechanisms of change in psychotherapy research. *Annu. Rev. Clin. Psychol.* **3**, 1–27 (2007).
71. Domhardt, M. et al. Mediators and mechanisms of change in internet- and mobile-based interventions for depression: a systematic review. *Clin. Psychol. Rev.* **83**, 101953 (2021).
72. Rad, M. S., Martingano, A. J. & Ginges, J. Toward a psychology of Homo sapiens: Making psychological science more representative of the human population. *Proc. Natl Acad. Sci. USA* **115**, 11401–11405 (2018).
73. Gone, J. P. A community-based treatment for Native American historical trauma: prospects for evidence-based practice. *J. Consult. Clin. Psychol.* **77**, 751–762 (2009).
74. Nagayama Hall, G. C., Yip, T. & Zárate, M. A. On becoming multicultural in a monocultural research world: a conceptual approach to studying ethnocultural diversity. *Am. Psychol.* **71**, 40–51 (2016).
75. Bernal, G. & Adames, C. Cultural adaptations: conceptual, ethical, contextual, and methodological issues for working with ethnocultural and majority-world populations. *Prev. Sci.* **18**, 681–688 (2017).
76. Castro, F. G., Barrera, M. & Holleran Steiker, L. K. Issues and challenges in the design of culturally adapted evidence-based interventions. *Ssrn* <https://doi.org/10.1146/annurev-clinpsy-033109-132032> (2010).
77. Warne, D. & Wescott, S. Social determinants of American Indian Nutritional Health. *Curr. Dev. Nutr.* <https://doi.org/10.1093/cdn/nzz054> (2019).
78. Abaza, H. & Marschollek, M. mHealth application areas and technology combinations. *Methods Inf. Med.* **56**, e105–e122 (2017).
79. Hall, C. S., Fottrell, E., Wilkinson, S. & Byass, P. Assessing the impact of mHealth interventions in low- and middle-income countries—what has been shown to work? *Glob. Health Action* **7**, 25606 (2014).
80. Kruse, C. et al. Barriers to the use of mobile health in improving health outcomes in developing countries: Systematic review. *J. Med. Internet Res.* **21**, 1–13 (2019).
81. Weisel, K. K. et al. Standalone smartphone apps for mental health—a systematic review and meta-analysis. *npj Digit. Med.* **2**, 1–10 (2019).
82. Bendig, E. et al. Internet-based interventions in chronic somatic disease. *Dtsch. Arztebl. Int.* **115**, 659–665 (2018).
83. Ebert, D. D. et al. Internet- and mobile-based psychological interventions: applications, efficacy, and potential for improving mental health. *Eur. Psychol.* **23**, 167–187 (2018).
84. Tsetsi, E. & Rains, S. A. Smartphone Internet access and use: extending the digital divide and usage gap. *Mob. Media Commun.* **5**, 239–255 (2017).
85. Baumeister, H., Reichler, L., Munzinger, M. & Lin, J. The impact of guidance on Internet-based mental health interventions - A systematic review. *Internet Inter.* **1**, 205–215 (2014).
86. Arsenijevic, J., Tummers, L. & Bosma, N. Adherence to electronic health tools among vulnerable groups: Systematic literature review and meta-analysis. *J. Med. Internet Res.* **22**, e11613 (2020).
87. Aranda-Jan, C. B., Mohutsiwa-Dibe, N. & Loukanova, S. Systematic review on what works, what does not work and why of implementation of mobile health (mHealth) projects in Africa. *BMC Public Health* **14**, 1–15 (2014).
88. James, D. C., Harville, C., Sears, C., Efunbumi, O. & Bondoc, I. Participation of African Americans in e-Health and m-Health studies: a systematic review. *Telemed. e-Health* **23**, 351–364 (2017).
89. Hwang, D. S. A., Lee, A., Song, J. M. & Han, H. R. Recruitment and retention strategies among racial and ethnic minorities in web-based intervention trials: retrospective qualitative analysis. *J. Med. Internet Res.* **23**, e23959 (2021).
90. Callier, S. & Fullerton, S. M. Diversity and inclusion in unregulated mHealth research: addressing the risks. *J. Law Med. Ethics* **48**, 115–121 (2020).
91. Druce, K. L., Dixon, W. G. & McBeth, J. Maximizing engagement in mobile health studies: lessons learned and future directions. *Rheum. Dis. Clin. North Am.* **45**, 159–172 (2019).
92. Moher, D., Liberati, A., Tetzlaff, J. & Altman, D. G. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *BMJ* **339**, 332–336 (2009).
93. Balci, S., Spanhel, K., Sander, L. & Baumeister, H. Protocol for a systematic review and meta-analysis of culturally adapted internet- And mobile-based health promotion interventions. *BMJ Open* <https://doi.org/10.1136/bmjopen-2020-037698> (2020).
94. Spanhel, K., Balci, S., Baumeister, H., Bengel, J. & Sander, L. B. Cultural adaptation of Internet- and mobile-based interventions for mental disorders: a systematic review protocol. *Syst. Rev.* **9**, 207 (2020).
95. Babineau, J. Product Review: Covidence (Systematic Review Software). *J. Can. Heal. Libr. Assoc./J. l'Association bibliothèques la St.é du Can.* **35**, 68 (2014).
96. Sterne, J. A. C. et al. RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ* <https://doi.org/10.1136/bmj.l4898> (2019).
97. Higgins JPT et al. (editors). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0. (The Cochrane Collaboration, 2011). Available at: www.handbook.cochrane.org.
98. Fritz, C. O., Morris, P. E. & Richler, J. J. Effect size estimates: current use, calculations, and interpretation. *J. Exp. Psychol. Gen.* **141**, 2–18 (2012).
99. Tufanaru, C., Munn, Z., Stephenson, M. & Aromataris, E. Fixed or random effects meta-analysis? Common methodological issues in systematic reviews of effectiveness. *Int. J. Evid. Based Health.* **13**, 196–207 (2015).
100. Balduzzi, S., Rucker, G. & Schwarzer, G. How to perform a meta-analysis with R: a practical tutorial. *Evid. Based Ment. Health* **22**, 153–160 (2019).
101. Harrer, M., Cuijpers, P. & Ebert, D. Doing meta-analysis in R. <https://doi.org/10.5281/ZENODO.2551803> (2019).
102. The Nordic Cochrane Centre. *Review Manager (RevMan)* (The Cochrane Collaboration, 2014).
103. Viechtbauer, W. Conducting meta-analyses in R with the metafor. *J. Stat. Softw.* **36**, 1–48 (2010).
104. Cohen, J. *Statistical Power Analysis for the Behavioural Science* 2nd edn. (Erlbaum Associate, 1988).
105. Higgins, J. P. T. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis. *Stat. Med.* <https://doi.org/10.1002/sim.1186> (2002).
106. Card, N. A. *Applied Meta-Analysis for Social Science Research*. (Guilford, 2011).
107. Bender, M. S. & Clark, M. J. Cultural adaptation for ethnic diversity: a review of obesity interventions for preschool children. *Calif. J. Health Promot* **9**, 40 (2011).
108. Barrera, M., Castro, F. G., Strycker, L. & Toobert, D. Cultural adaptations of behavioral health interventions: a progress report. *J. Consult. Clin. Psychol.* **81**, 196–205 (2013).
109. Castro, Y. et al. Adaptation of a counseling intervention to address multiple cancer risk factors among overweight/obese latino smokers. *Heal. Educ. Behav.* **42**, 65–72 (2015).
110. Philis-Tsimikas, A. et al. Improvement in diabetes care of underinsured patients enrolled in Project Dulce: A community-based, culturally appropriate, nurse case management and peer education diabetes care model. *Diabetes Care* **27**, 1110–1115 (2004).

ACKNOWLEDGEMENTS

The authors wish to thank all contacted authors who provided the missing information. We would like to thank student assistants Fatma Kahir, Karamdeep Khinda, and Felicitas Feldhahn for their support in screening and data extraction. S.B. receives a scholarship granted by the Ministry of National Education in Turkey, K.S. is supported by the German Academic Scholarship Foundation. Open Access funding was provided by the University of Ulm. The funders had no role in study design, data collection, and analysis, decision to publish, or preparation of the manuscript.

AUTHOR CONTRIBUTIONS

S.B., K.S., L.S., and H.B. conceived the study design. S.B., K.S., and L.S. developed the search strategy, study selection, and extraction. S.B. wrote the draft of the manuscript. All authors read, provided feedback, and approved the final version. S.B. is the guarantor of the review.

FUNDING

Open Access funding enabled and organized by Projekt DEAL.

COMPETING INTERESTS

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41746-022-00569-x>.

Correspondence and requests for materials should be addressed to Sumeyye Balci.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022